

Seed orchad genetic

Origin	<i>Abies nordmanniana</i>		<i>Abies bornmülleriana</i>	
	Ambrolauri	Tversted	North Caucasus	Bornmülleriana
1. gen.	(1995) Skibelund FP.266	(1995) Silkeborg Nordskov FP.259	(1964) Vallo FP.620	(1995) Kongsøre FP.267
	(1996) Sønderskovgaard FP.246	(2001) Mosemarkskov FP.251	(1997) Vargårde FP.254	
			(2016) Kongeå FP.1021 (orig. Apsheronsk)	
			(2025) Rostock FP.1024 (orig. Apsheronsk)	
			(2025) Hillerød FP.1025 (orig. Apsheronsk)	
1.5 gen.	(2012) Mikkelborg FP.1004 (orig. FP.266)	(1988) Bredal FP.1100		(2014) Ry FP.1005 (orig. FP.267)
		(1997) Skelhusmarken FP.272 (orig. Boller & Tversted)		(2019) Gl. Ry FP.1006 (orig. Kongsøre FP.267)
		(2007) Lundbygaard FP.1000		
		(2008) Stursbøl plantage FP.273		
		(2008) Smørhaverne FP.1101		
		(2012) Kalby plantage FP.670		
		(2012) Lundbygaard FP.1001		
		(2012) Dramstrup FP.1002		
		(2012) Aars FP.1003		
		(2015) Kalby plantage FP.690		
2. gen.	(2024) Gl. Kirstineberg FP.1015 (orig. FP.266)			(202x) Ry FP1023 1006 (orig. Kongsøre FP.267)
	(2024) Gl. Kirstineberg FP.1016 (orig. FP.266)			
	(2024) Gl. Kirstineberg FP.1017 (orig. FP.266)			
	(202x) Hillerød FP.1022 (orig. FP.266)			